

# Quarkonium physics at a fixed-target experiment with the proton and lead LHC beams

**Jean-Philippe Lansberg**  
IPNO, Paris-Sud XI U.

**Brookhaven Summer Program, Quarkonium Production in  
Elementary and Heavy Ion Collisions**

June 17, 2011

Brookhaven National Laboratory, USA

with F. Fleuret (LLR), S.J. Brodsky (SLAC), ...

# Part I

## A fixed-target experiment using the LHC beam(s): generalities

# A Fixed Target Experiment

## Generalities

- $pp$  or  $pA$  with a 7 TeV  $p$  beam :  $\sqrt{s} \simeq 115 \text{ GeV}$  (+Fermi motion for  $pA$ )

# A Fixed Target Experiment

## Generalities

- $pp$  or  $pA$  with a 7 TeV  $p$  beam :  $\sqrt{s} \simeq 115 \text{ GeV}$  (+Fermi motion for  $pA$ )
- Same ballpark as electron-ion colliders → **complementary**

# A Fixed Target Experiment

## Generalities

- $pp$  or  $pA$  with a 7 TeV  $p$  beam :  $\sqrt{s} \simeq 115 \text{ GeV}$  (+Fermi motion for  $pA$ )
- Same ballpark as electron-ion colliders → **complementary**
- For  $pA$ , a Fermi motion of 0.2 GeV would induce a spread of 10 % of  $\sqrt{s}$

S.Fredriksson, NPB 94 (1975) 337

# A Fixed Target Experiment

## Generalities

- $pp$  or  $pA$  with a 7 TeV  $p$  beam :  $\sqrt{s} \simeq 115 \text{ GeV}$  (+Fermi motion for  $pA$ )
- Same ballpark as electron-ion colliders  $\rightarrow$  **complementary**
- For  $pA$ , a Fermi motion of 0.2 GeV would induce a spread of 10 % of  $\sqrt{s}$
- The beam may be extracted using “Strong cristalline field”

S.Fredriksson, NPB 94 (1975) 337

E. Huggerhøj, U.I Huggerhøj, NIM B 234 (2005) 31, Rev. Mod. Phys. 77 (2005) 1131 (+ next page)

# A Fixed Target Experiment

## Generalities

- $pp$  or  $pA$  with a 7 TeV  $p$  beam :  $\sqrt{s} \simeq 115 \text{ GeV}$  (+Fermi motion for  $pA$ )
- Same ballpark as electron-ion colliders  $\rightarrow$  **complementary**
- For  $pA$ , a Fermi motion of 0.2 GeV would induce a spread of 10 % of  $\sqrt{s}$

S.Fredriksson, NPB 94 (1975) 337

- The beam may be extracted using “Strong cristalline field”

E. Huggerhøj, U.I Huggerhøj, NIM B 234 (2005) 31, Rev. Mod. Phys. 77 (2005) 1131 (+ next page)

- Expected luminosities with  $5 \times 10^8 \text{ p/s}$  extracted (1cm-long target)

Target	$\rho \text{ (g.cm}^{-3}\text{)}$	A	$\mathcal{L} \text{ (}\mu\text{b}^{-1}\text{.s}^{-1}\text{)}$	$\mathcal{L} \text{ (pb}^{-1}\text{.y}^{-1}\text{)}$
Liq. H <sub>2</sub>	0.07	1	21	210
Liq. D <sub>2</sub>	0.16	2	24	240
Be	1.85	9	60	600
Cu	8.96	64	40	400
W	19.1	185	30	300
Pb	11.35	207	16	160

(preliminary !)

# A Fixed Target Experiment

## Generalities

- $pp$  or  $pA$  with a 7 TeV  $p$  beam :  $\sqrt{s} \simeq 115 \text{ GeV}$  (+Fermi motion for  $pA$ )
- Same ballpark as electron-ion colliders  $\rightarrow$  **complementary**
- For  $pA$ , a Fermi motion of 0.2 GeV would induce a spread of 10 % of  $\sqrt{s}$

S.Fredriksson, NPB 94 (1975) 337

- The beam may be extracted using “Strong cristalline field”  
E. Huggerhøj, U.I Huggerhøj, NIM B 234 (2005) 31, Rev. Mod. Phys. 77 (2005) 1131 (+ next page)
- Expected luminosities with  $5 \times 10^8 \text{ p/s}$  extracted (1cm-long target)

Target	$\rho \text{ (g.cm}^{-3}\text{)}$	A	$\mathcal{L} \text{ (}\mu\text{b}^{-1}.\text{s}^{-1}\text{)}$	$\mathcal{L} \text{ (pb}^{-1}.\text{y}^{-1}\text{)}$
Liq. $\text{H}_2$	0.07	1	21	210
Liq. $\text{D}_2$	0.16	2	24	240
Be	1.85	9	60	600
Cu	8.96	64	40	400
W	19.1	185	30	300
Pb	11.35	207	16	160

(preliminary !)

- Using **NA51**-like 1.2m-long liquid  $\text{H}_2$  &  $\text{D}_2$  targets,  $\mathcal{L}_{\text{H}_2/\text{D}_2} \simeq 20 \text{ fb}^{-1} \text{y}^{-1}$



# A Fixed Target Experiment

## Generalities

- $pp$  or  $pA$  with a 7 TeV  $p$  beam :  $\sqrt{s} \simeq 115 \text{ GeV}$  (+Fermi motion for  $pA$ )
- Same ballpark as electron-ion colliders  $\rightarrow$  **complementary**
- For  $pA$ , a Fermi motion of 0.2 GeV would induce a spread of 10 % of  $\sqrt{s}$

S.Fredriksson, NPB 94 (1975) 337

- The beam may be extracted using “Strong cristalline field”  
E. Huggerhøj, U.I Huggerhøj, NIM B 234 (2005) 31, Rev. Mod. Phys. 77 (2005) 1131 (+ next page)
- Expected luminosities with  $5 \times 10^8 \text{ p/s}$  extracted (1cm-long target)

Target	$\rho \text{ (g.cm}^{-3}\text{)}$	A	$\mathcal{L} \text{ (}\mu\text{b}^{-1}.\text{s}^{-1}\text{)}$	$\mathcal{L} \text{ (pb}^{-1}.\text{y}^{-1}\text{)}$
Liq. $\text{H}_2$	0.07	1	21	210
Liq. $\text{D}_2$	0.16	2	24	240
Be	1.85	9	60	600
Cu	8.96	64	40	400
W	19.1	185	30	300
Pb	11.35	207	16	160

(preliminary !)

- Using **NA51**-like 1.2m-long liquid  $\text{H}_2$  &  $\text{D}_2$  targets,  $\mathcal{L}_{\text{H}_2/\text{D}_2} \simeq 20 \text{ fb}^{-1} \text{y}^{-1}$
- For comparison, PHENIX recorded lumi for  
Run9 pp at 200 GeV:  $16 \text{ pb}^{-1}$  & Run8 dAu at 200 GeV :  $0.08 \text{ pb}^{-1}$

# A Fixed Target Experiment

## Generalities

- *Pbp* or *PbA* with a 2.75 TeV Pb beam :  $\sqrt{s} \simeq 72 \text{ GeV}$

# A Fixed Target Experiment

## Generalities

- *Pbp* or *PbA* with a 2.75 TeV Pb beam :  $\sqrt{s} \simeq 72 \text{ GeV}$
- Cristal channeling is also possible (to extract a few per cent of the beam)

# A Fixed Target Experiment

## Generalities

- *Pbp* or *PbA* with a 2.75 TeV Pb beam :  $\sqrt{s} \simeq 72 \text{ GeV}$
- Cristal channeling is also possible (to extract a few per cent of the beam)
- Requires cristals highly resistant to radiations: progress with diamonds

P. Ballin *et al.*, NIMB 267 (2009) 2952

# A Fixed Target Experiment

## Generalities

- *Pbp* or *PbA* with a 2.75 TeV Pb beam :  $\sqrt{s} \simeq 72 \text{ GeV}$
- Cristal channeling is also possible (to extract a few per cent of the beam)
- Requires cristals highly resistant to radiations: progress with diamonds

P. Ballin *et al.*, NIMB 267 (2009) 2952

- Expected luminosities with  $7 \times 10^5 \text{ Pb/s}$  extracted (1cm-long target)

Target	$\rho \text{ (g.cm}^{-3}\text{)}$	A	$\mathcal{L} \text{ (mb}^{-1}\text{s}^{-1}\text{)} = f \mathcal{L} \text{ (nb}^{-1}\text{.yr}^{-1}\text{)}$
Liq. H <sub>2</sub>	0.07	1	28
Liq. D <sub>2</sub>	0.16	2	34
Be	1.85	9	84
Cu	8.96	64	56
W	19.1	185	42
Pb	11.35	207	22

(Preliminary !)

# A Fixed Target Experiment

## Generalities

- *Pbp* or *PbA* with a 2.75 TeV Pb beam :  $\sqrt{s} \simeq 72 \text{ GeV}$
- Cristal channeling is also possible (to extract a few per cent of the beam)
- Requires cristals highly resistant to radiations: progress with diamonds

P. Ballin *et al.*, NIMB 267 (2009) 2952

- Expected luminosities with  $7 \times 10^5 \text{ Pb/s}$  extracted (1cm-long target)

Target	$\rho \text{ (g.cm}^{-3}\text{)}$	A	$\mathcal{L} \text{ (mb}^{-1}\text{s}^{-1}\text{)} = f \mathcal{L} \text{ (nb}^{-1}\text{.yr}^{-1}\text{)}$
Liq. H <sub>2</sub>	0.07	1	28
Liq. D <sub>2</sub>	0.16	2	34
Be	1.85	9	84
Cu	8.96	64	56
W	19.1	185	42
Pb	11.35	207	22

(Preliminary !)

- For comparison, Phenix recorded lumi for Run10  
AuAu at 200 GeV:  $1.3 \text{ nb}^{-1}$  & AuAu at 62 GeV:  $0.11 \text{ nb}^{-1}$


# Beam extraction

## • Beam extraction @ LHC

... there are extremely promising possibilities to extract 7 TeV protons from the circulating beam by means of a bent crystal.

... The idea is to put a bent, single crystal of either Si or Ge (W would perform slightly better but needs substantial improvements in crystal quality) at a distance of  $\simeq 7\sigma$  to the beam where it can intercept and deflect part of the beam halo by an angle similar to the one the foreseen dump kicking system will apply to the circulating beam.

... ions with the same momentum per charge as protons are deflected in a crystal with similar efficiencies



If the crystal is positioned at the kicking section, the whole dump system can be used for slow extraction of parts of the beam halo, the particles that are anyway lost subsequently at collimators.

## Part II

### AFTER as a quarkonium observatory in $pp$

(constraining the glue at large  $x$  in the proton)



# Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**

# Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
  - in photo/lepto production (DIS)
  - but also in  $g - g$ -fusion process

# Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
  - in photo/lepto production (DIS)
  - but also in  $g - g$ -fusion process
  - mainly because of the presence of a natural “hard” scale:  $m_Q$
  - and the good detectability of a dimuon pair

# Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
  - in photo/lepto production (DIS)
  - but also in  $g - g$ -fusion process
  - mainly because of the presence of a natural “hard” scale:  $m_Q$
  - and the good detectability of a dimuon pair

PHYSICAL REVIEW D

VOLUME 37, NUMBER 5

1 MARCH 1988

## Structure-function analysis and $\psi$ , jet, $W$ , and $Z$ production: Determining the gluon distribution

A. D. Martin

*Department of Physics, University of Durham, Durham, England*

R. G. Roberts

*Rutherford Appleton Laboratory, Didcot, Oxon, England*

W. J. Stirling

*Department of Physics, University of Durham, Durham, England*

(Received 27 July 1987)

We perform a next-to-leading-order structure-function analysis of deep-inelastic  $\mu N$  and  $\nu N$  scattering data and find acceptable fits for a range of input gluon distributions. We show three equally acceptable sets of parton distributions which correspond to gluon distributions which are (1) “soft,” (2) “hard,” and (3) which behave as  $xG(x) \sim 1/\sqrt{x}$  at small  $x$ .  $J/\psi$  and prompt photon hadroproduction data are used to discriminate between the three sets. Set 1, with the “soft”-gluon distribution, is favored.  $W$ ,  $Z$ , and jet production data from the CERN collider are well described but do not distinguish between the sets of structure functions. The precision of the predictions for  $\sigma_W$  and  $\sigma_Z$  allow the collider measurements to yield information on the number of light neutrinos and the mass of the top quark. Finally we discuss how the gluon distribution at very small  $x$  may be directly measured at DESY HERA.

# Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
  - in photo/lepto production (DIS)
  - but also in  $g - g$ -fusion process
  - mainly because of the presence of a natural “hard” scale:  $m_Q$
  - and the good detectability of a dimuon pair

Z. Phys. C – Particles and Fields 38, 473–478 (1988)

## $J/\psi$ Production at large transverse momentum at hadron colliders

E.W.N. Glover<sup>1</sup>\*, A.D. Martin<sup>2</sup>, W.J. Stirling<sup>2</sup>

<sup>1</sup> Cavendish Laboratory, University of Cambridge, Cambridge, CB3 0HE, England

<sup>2</sup> Physics Department, University of Durham, Durham, DH1 3LE, England

Received 7 October 1987

**Abstract.** We calculate  $J/\psi$  hadroproduction and emphasize the importance of the  $J/\psi$  signal as a measure of  $b\bar{b}$  production via the decay  $B \rightarrow \psi X$  and of the gluon structure function at low  $x$  via  $\chi$  hadroproduction followed by  $\chi \rightarrow \psi \gamma$  decay. We compare with UA1 data and data at ISR energies and make predictions for  $\psi$  production at TEVATRON energies.

# Need for a quarkonium observatory

- Many **hopes** were put in **quarkonium studies** to extract **gluon PDF**
  - in photo/lepto production (DIS)
  - but also in  $g - g$ -fusion process
  - mainly because of the presence of a natural “hard” scale:  $m_Q$
  - and the good detectability of a dimuon pair

PHYSICAL REVIEW D

VOLUME 48, NUMBER 11

1 DECEMBER 1993

## $\psi$ production in $\bar{p}N$ and $\pi^- N$ interactions at 125 GeV/c and a determination of the gluon structure functions of the $\bar{p}$ and the $\pi^-$

C. Akerlof,<sup>4</sup> H. Areti,<sup>3,\*</sup> M. Binkley,<sup>2</sup> S. Conetti,<sup>3,†</sup> B. Cox,<sup>2,†</sup> J. Enagonio,<sup>2</sup>  
He Mao,<sup>5</sup> C. Hojvat,<sup>2</sup> D. Judd,<sup>2,‡</sup> S. Katsanevas,<sup>1</sup> R. D. Kephart,<sup>2</sup> C. Kourkouvelis,<sup>1</sup> P. Kraushaar,<sup>4,§</sup>  
P. Lebrun,<sup>3,\*</sup> P. K. Malhotra,<sup>2,||</sup> A. Markou,<sup>1</sup> P. O. Mazur,<sup>2</sup> D. Nitz,<sup>4</sup> L. K. Resvanis,<sup>1</sup> D. Ryan,<sup>3</sup>  
T. Ryan,<sup>3,¶</sup> W. Schappert,<sup>3,\*\*</sup> D. G. Stairs,<sup>3</sup> R. Thun,<sup>4</sup> F. Turkot,<sup>2</sup> S. Tzamarias,<sup>1,||</sup> G. Voulgaris,<sup>1</sup>  
R. L. Wagner,<sup>2</sup> D. E. Wagoner,<sup>2,‡</sup> W. Yang,<sup>2</sup> and Zhang Nai-jian<sup>5</sup>

(E537 Collaboration)

<sup>1</sup>University of Athens, Athens, Greece

<sup>2</sup>Fermi National Accelerator Laboratory, Batavia, Illinois 60510

<sup>3</sup>McGill University, Montreal, Quebec, Canada H3A 2T8

<sup>4</sup>University of Michigan, Ann Arbor, Michigan 48109

<sup>5</sup>Shandong University, Jinan, People's Republic of China

(Received 9 February 1993)

We have measured the cross section for production of  $\psi$  and  $\psi'$  in  $\bar{p}$  and  $\pi^-$  interactions with Be, Cu, and W targets in experiment E537 at Fermilab. The measurements were performed at 125 GeV/c using a forward dimuon spectrometer in a closed geometry configuration. The gluon structure functions of the  $\bar{p}$  and  $\pi^-$  have been extracted from the measured  $d\sigma/dx_F$  spectra of the produced  $\psi$ 's. From the  $\bar{p}W$  data we obtain, for  $\bar{p}$ ,  $xG(x) = (2.15 \pm 0.7)[1-x]^{(6.83 \pm 0.5)} [1 + (5.85 \pm 0.95)x]$ . In the  $\pi^-$  case, we obtain, from the W and the Be data separately,  $xG(x) = (1.49 \pm 0.03)[1-x]^{(1.98 \pm 0.06)}$  (for  $\pi^-W$ ),  $xG(x) = (1.10 \pm 0.10)[1-x]^{(1.20 \pm 0.20)}$  (for  $\pi^-Be$ ).

# Need for a quarkonium observatory

- Big theoretical complications

# Need for a quarkonium observatory

- Big theoretical complications
  - Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )



# Need for a quarkonium observatory

- Big theoretical complications

- Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )
- Naive application of pQCD (CSM) fails to describe  $d\sigma/dP_T$  at LO  
(after all, it is at large  $P_T$  that it would be safer to extract reliable info)

# Need for a quarkonium observatory

- **Big theoretical complications**

- Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )
- Naive application of pQCD (CSM) fails to describe  $d\sigma/dP_T$  at LO  
(after all, it is at large  $P_T$  that it would be safer to extract reliable info)
- Different competing models: CSM, NRQCD-COM, CEM,  $k_T$  fact.

# Need for a quarkonium observatory

- **Big theoretical complications**

- Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )
- Naive application of pQCD (CSM) fails to describe  $d\sigma/dP_T$  at LO  
(after all, it is at large  $P_T$  that it would be safer to extract reliable info)
- Different competing models: CSM, NRQCD-COM, CEM,  $k_T$  fact.
- At larger  $\sqrt{s}$  (or  $P_T$ ), the  $B$  enter the game

# Need for a quarkonium observatory

- **Big theoretical complications**

- Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )
- Naive application of pQCD (CSM) fails to describe  $d\sigma/dP_T$  at LO  
(after all, it is at large  $P_T$  that it would be safer to extract reliable info)
- Different competing models: CSM, NRQCD-COM, CEM,  $k_T$  fact.
- At larger  $\sqrt{s}$  (or  $P_T$ ), the  $B$  enter the game
- All this calls for very involved theoretical computations or experimental measurements

# Need for a quarkonium observatory

- **Big theoretical complications**

- Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )
- Naive application of pQCD (CSM) fails to describe  $d\sigma/dP_T$  at LO  
(after all, it is at large  $P_T$  that it would be safer to extract reliable info)
- Different competing models: CSM, NRQCD-COM, CEM,  $k_T$  fact.
- At larger  $\sqrt{s}$  (or  $P_T$ ), the  $B$  enter the game

- All this calls for very involved theoretical computations or experimental measurements
- **Even at low  $P_T$ , things are not easy**

# Need for a quarkonium observatory

- **Big theoretical complications**

- Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )
- Naive application of pQCD (CSM) fails to describe  $d\sigma/dP_T$  at LO  
(after all, it is at large  $P_T$  that it would be safer to extract reliable info)
- Different competing models: CSM, NRQCD-COM, CEM,  $k_T$  fact.
- At larger  $\sqrt{s}$  (or  $P_T$ ), the  $B$  enter the game

- All this calls for very involved theoretical computations or experimental measurements

- **Even at low  $P_T$ , things are not easy**

- Specific difficulties to measure the  $\chi_{c,b}$
- Reduced acceptance at the LHC (CMS, ATLAS)

# Need for a quarkonium observatory

- **Big theoretical complications**

- Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )
- Naive application of pQCD (CSM) fails to describe  $d\sigma/dP_T$  at LO  
(after all, it is at large  $P_T$  that it would be safer to extract reliable info)
- Different competing models: CSM, NRQCD-COM, CEM,  $k_T$  fact.
- At larger  $\sqrt{s}$  (or  $P_T$ ), the  $B$  enter the game

- All this calls for very involved theoretical computations or experimental measurements

- **Even at low  $P_T$ , things are not easy**

- Specific difficulties to measure the  $\chi_{c,b}$
- Reduced acceptance at the LHC (CMS, ATLAS)
- Very large theoretical uncertainty (mass,  $\alpha_s(\mu_R)$ )

# Need for a quarkonium observatory

- **Big theoretical complications**

- Direct production is not dominant (neither indirect –via  $\chi_{Q-}$ )
- Naive application of pQCD (CSM) fails to describe  $d\sigma/dP_T$  at LO  
(after all, it is at large  $P_T$  that it would be safer to extract reliable info)
- Different competing models: CSM, NRQCD-COM, CEM,  $k_T$  fact.
- At larger  $\sqrt{s}$  (or  $P_T$ ), the  $B$  enter the game

- All this calls for very involved theoretical computations or experimental measurements

- **Even at low  $P_T$ , things are not easy**

- Specific difficulties to measure the  $\chi_{c,b}$
- Reduced acceptance at the LHC (CMS, ATLAS)
- Very large theoretical uncertainty (mass,  $\alpha_s(\mu_R)$ )
- **Yet, very sensitive on  $g(x, Q^2)$**  where it is not well known



# Need for a quarkonium observatory

When hadroproduction casts doubt on lepto/photo-production

# Need for a quarkonium observatory

When hadroproduction casts doubt on lepto/photo-production

Volume 191, number 1,2

PHYSICS LETTERS B

4 June 1987

## INELASTIC LEPTOPRODUCTION OF $J/\psi$ AS A PROBE OF THE SMALL- $x$ BEHAVIOUR OF THE GLUON STRUCTURE FUNCTION

A.D. MARTIN, C.-K. NG and W.J. STIRLING

*Department of Physics, University of Durham, Durham DH1 3LE, UK*

Received 19 February 1987

The differential cross section  $d\sigma/dx$  for the inelastic photoproduction of  $J/\psi$  is predicted, via the subprocess  $\gamma g \rightarrow \psi g$ , to be sharply peaked at a small value,  $x_{\text{peak}}$ , of  $x$  and, even more remarkably, the integrated  $\gamma N \rightarrow \psi X$  cross section is, up to a calculable numerical constant, essentially the proton-gluon distribution  $xG(x)$  at  $x \simeq x_{\text{peak}}$ . Cross section measurements at HERA may thus provide a direct determination of  $G(x)$  for  $x \sim 10^{-3}$ . Inelastic  $J/\psi$  events arising from  $b\bar{b}$  production are also studied.

# Need for a quarkonium observatory

## When hadroproduction casts doubt on lepto/photo-production

Volume 258, number 3,4

PHYSICS LETTERS B

11 April 1991

### Inelastic $J/\psi$ production in deep inelastic scattering from hydrogen and deuterium and the gluon distribution of free nucleons

New Muon Collaboration (NMC)


Received 10 November 1990

We present results on inelastic  $J/\psi$  production from muon interactions with hydrogen and deuterium at an incident muon energy of 280 GeV. The measured cross section ratio per nucleon for muon-induced  $J/\psi$  production in deuterium and hydrogen was found to be  $R(D_2/H_2) = 1.01 \pm 0.15$ . The colour singlet model is shown to provide a good description of the observed differential cross section apart from a normalisation factor. The comparison between the observed cross section and the colour singlet model prediction allows the extraction of the gluon structure function  $G(x)$  of the nucleon. The momentum fraction  $x$  of the nucleon carried by the gluon is measured in the range of  $x = [0.02, 0.30]$ . The normalised gluon distribution of free nucleons thus found can be parametrised as  $xG(x) = \frac{1}{2}(\eta + 1)(1 - x)^\eta$ , with  $\eta = 5.1 \pm 0.9$  (stat.).

# Need for a quarkonium observatory

When hadroproduction casts doubt on lepto/photo-production

Even though, the LO CSM fails to describe the data. the NLO does a good job  
(we are in 1996 !)




M.Kramer Nucl.Phys.B459:3 1996  
H1,EPJC 25, 2,2002; ZEUS, EPJC 27, 173, 2003

# Need for a quarkonium observatory

When hadroproduction casts doubt on lepto/photo-production

Even though, the LO CSM fails to describe the data. the NLO does a good job  
(we are in 1996 !)




M.Kramer Nucl.Phys.B459:3 1996  
H1,EPJC 25, 2,2002; ZEUS, EPJC 27, 173, 2003

# Need for a quarkonium observatory

When hadroproduction casts doubt on lepto/photo-production

Even though, the LO CSM fails to describe the data. the NLO does a good job  
(we are in 1996 !)




M.Kramer Nucl.Phys.B459:3 1996  
H1,EPJC 25, 2,2002; ZEUS, EPJC 27, 173, 2003

- Yet, since then,  $J/\psi$  data are **not included in global fits anymore**

# Need for a quarkonium observatory

When hadroproduction casts doubt on lepto/photo-production

Even though, the LO CSM fails to describe the data. the NLO does a good job  
(we are in 1996 !)




M.Kramer Nucl.Phys.B459:3 1996  
H1,EPJC 25, 2,2002; ZEUS, EPJC 27, 173, 2003

- Yet, since then,  $J/\psi$  data are **not included in global fits anymore**
- Is it because other means are more precise ?
- or the  $J/\psi$  probe is not reliable as long as pp data are not understood ?

# Need for a quarkonium observatory

When hadroproduction casts doubt on lepto/photo-production

Even though, the LO CSM fails to describe the data. the NLO does a good job  
(we are in 1996 !)



M.Kramer Nucl.Phys.B459:3 1996  
H1,EPJC 25, 2,2002; ZEUS, EPJC 27, 173, 2003

- Yet, since then,  $J/\psi$  data are **not included in global fits anymore**
- Is it because other means are more precise ?
- or the  $J/\psi$  probe is not reliable as long as pp data are not understood ?
- This issue needs also to be addressed when discussing **EIC**



# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield (  $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  (  $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)

# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield (  $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  (  $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)
- $\chi_{c,b}$  production is **badly known**, even **worse for the  $\eta_c$**

# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield (  $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  (  $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)
- $\chi_{c,b}$  production is **badly known**, even **worse for the  $\eta_c$**
- The latter are **potentially better probes** of glue in  $pp$

# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield (  $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  (  $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)
- $\chi_{c,b}$  production is **badly known**, even **worse for the  $\eta_c$**
- The latter are **potentially better probes** of glue in  $pp$
- LO processes are  $gg \rightarrow \begin{cases} \chi_{c,b,2} \\ \eta_{c,b} \end{cases}$

# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield ( $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  ( $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)
- $\chi_{c,b}$  production is **badly known**, even **worse for the  $\eta_c$**
- The latter are **potentially better probes** of glue in  $pp$
- LO processes are  $gg \rightarrow \begin{cases} \chi_{c,b,2} \\ \eta_{c,b} \end{cases}$
- For that, we need

# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield ( $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  ( $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)
- $\chi_{c,b}$  production is **badly known**, even **worse for the  $\eta_c$**
- The latter are **potentially better probes** of glue in  $pp$
- LO processes are  $gg \rightarrow \begin{cases} \chi_{c,b,2} \\ \eta_{c,b} \end{cases}$
- For that, we need
  - **high stats**  
→ wide acceptance (also help not to bias 1D polarisation analyses)

# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield ( $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  ( $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)
- $\chi_{c,b}$  production is **badly known**, even **worse for the  $\eta_c$**
- The latter are **potentially better probes** of glue in  $pp$
- LO processes are  $gg \rightarrow \begin{cases} \chi_{c,b,2} \\ \eta_{c,b} \end{cases}$
- For that, we need
  - **high stats**  
→ wide acceptance (also help not to bias 1D polarisation analyses)
  - a vertex detector

# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield ( $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  ( $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)
- $\chi_{c,b}$  production is **badly known**, even **worse for the  $\eta_c$**
- The latter are **potentially better probes** of glue in  $pp$
- LO processes are  $gg \rightarrow \begin{cases} \chi_{c,b,2} \\ \eta_{c,b} \end{cases}$
- For that, we need
  - **high stats**  
→ wide acceptance (also help not to bias 1D polarisation analyses)
  - a vertex detector
  - state-of-the-art calorimetry for  $\gamma$  ( $\chi_Q \rightarrow {}^3S_1 + \gamma$ ,  $\eta_c \rightarrow \gamma\gamma$ )



# Need for a quarkonium observatory

- To put an end to production controversies (since 1995 !), we need
  - a study of **direct**  $J/\psi$  yield ( $\chi_c$  only measured in  $pp$  by CDF and PHENIX)
  - a study of **direct**  $\Upsilon(nS)$  ( $\chi_b$  only measured in  $pp$  by CDF (1 point))
  - a study of the polarisation of **direct** yields  
(at least in 2 frames or 2D distrib.)
  - + probably associated production (see later)
- $\chi_{c,b}$  production is **badly known**, even **worse for the  $\eta_c$**
- The latter are **potentially better probes** of glue in  $pp$
- LO processes are  $gg \rightarrow \begin{cases} \chi_{c,b,2} \\ \eta_{c,b} \end{cases}$
- For that, we need
  - **high stats**  
→ wide acceptance (also help not to bias 1D polarisation analyses)
  - a vertex detector
  - state-of-the-art calorimetry for  $\gamma$  ( $\chi_Q \rightarrow {}^3S_1 + \gamma$ ,  $\eta_c \rightarrow \gamma\gamma$ )
  - adapted triggers (Big issue for CMS and ATLAS)

# A Fixed Target Experiment: A quarkonium observatory

- Interpolating the world data set:

Target	$N_{J/\psi} \text{ (}\gamma^{-1}\text{)}$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_{\Upsilon} \text{ (}\gamma^{-1}\text{)}$ <small><math>N_{\Upsilon} = A \mathcal{L} \sigma_{\Upsilon}</math></small>
(with branching and per unit of rapidity)		
Liq. H <sup>2</sup> (1m)	0.6 10 <sup>9</sup>	10 <sup>6</sup>
Liq. D <sup>2</sup>	1.5 10 <sup>9</sup>	23 10 <sup>5</sup>
Be	0.2 10 <sup>9</sup>	2.7 10 <sup>5</sup>
Cu	0.8 10 <sup>9</sup>	13 10 <sup>5</sup>
W	1.7 10 <sup>9</sup>	27 10 <sup>5</sup>
Pb	1. 10 <sup>9</sup>	16 10 <sup>5</sup>

## A Fixed Target Experiment: A quarkonium observatory

- Interpolating the world data set:
- Rates expected at RHIC in 2011:  
 $J/\psi$ :  $10^6$  in  $pp$ ,  $\Upsilon$ :  $10^4$  in  $pp$

Target	$N_{J/\psi} \text{ (}\gamma^{-1}\text{)}$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_{\Upsilon} \text{ (}\gamma^{-1}\text{)}$ <small><math>N_{\Upsilon} = A \mathcal{L} \sigma_{\Upsilon}</math></small>
(with branching and per unit of rapidity)		
Liq. H <sup>2</sup> (1m)	0.6 $10^9$	$10^6$
Liq. D <sup>2</sup>	1.5 $10^9$	23 $10^5$
Be	0.2 $10^9$	2.7 $10^5$
Cu	0.8 $10^9$	13 $10^5$
W	1.7 $10^9$	27 $10^5$
Pb	1. $10^9$	16 $10^5$

## A Fixed Target Experiment: A quarkonium observatory

- Interpolating the world data set:
- Rates expected at RHIC in 2011:  
 $J/\psi$ :  $10^6$  in  $pp$ ,  $\Upsilon$ :  $10^4$  in  $pp$
- 2-3 orders of magnitude higher here  
(RHIC yields are much lower in dAu compared to pA here)

Target	$N_{J/\psi} \text{ (}\gamma^{-1}\text{)}$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_{\Upsilon} \text{ (}\gamma^{-1}\text{)}$ <small><math>N_{\Upsilon} = A \mathcal{L} \sigma_{\Upsilon}</math></small>
(with branching and per unit of rapidity)		
Liq. H <sup>2</sup> (1m)	0.6 10 <sup>9</sup>	10 <sup>6</sup>
Liq. D <sup>2</sup>	1.5 10 <sup>9</sup>	23 10 <sup>5</sup>
Be	0.2 10 <sup>9</sup>	2.7 10 <sup>5</sup>
Cu	0.8 10 <sup>9</sup>	13 10 <sup>5</sup>
W	1.7 10 <sup>9</sup>	27 10 <sup>5</sup>
Pb	1. 10 <sup>9</sup>	16 10 <sup>5</sup>

## A Fixed Target Experiment: A quarkonium observatory

- Interpolating the world data set:
- Rates expected at RHIC in 2011:  
 $J/\psi$ :  $10^6$  in  $pp$ ,  $\Upsilon$ :  $10^4$  in  $pp$
- 2-3 orders of magnitude higher here  
(RHIC yields are much lower in dAu compared to pA here)
- Numbers are for only one unit of  $y$  about 0

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{J/\psi}</math></small>	$N_{\Upsilon} (y^{-1})$ <small><math>N_{\Upsilon} = A \mathcal{L} \sigma_{\Upsilon}</math></small>
(with branching and per unit of rapidity)		
Liq. H <sup>2</sup> (1m)	0.6 $10^9$	$10^6$
Liq. D <sup>2</sup>	1.5 $10^9$	23 $10^5$
Be	0.2 $10^9$	2.7 $10^5$
Cu	0.8 $10^9$	13 $10^5$
W	1.7 $10^9$	27 $10^5$
Pb	1. $10^9$	16 $10^5$

## A Fixed Target Experiment: A quarkonium observatory


- Interpolating the world data set:
- Rates expected at RHIC in 2011:  
 $J/\psi$ :  $10^6$  in  $pp$ ,  $\Upsilon$ :  $10^4$  in  $pp$
- 2-3 orders of magnitude higher here  
(RHIC yields are much lower in dAu compared to pA here)
- Numbers are for only one unit of  $y$  about 0
- Unique access in the backward region

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{J/\psi}</math></small>	$N_{\Upsilon} (y^{-1})$ <small><math>N_{\Upsilon} = A \mathcal{L} \sigma_{\Upsilon}</math></small>
(with branching and per unit of rapidity)		
Liq. H <sup>2</sup> (1m)	0.6 $10^9$	$10^6$
Liq. D <sup>2</sup>	1.5 $10^9$	23 $10^5$
Be	0.2 $10^9$	2.7 $10^5$
Cu	0.8 $10^9$	13 $10^5$
W	1.7 $10^9$	27 $10^5$
Pb	1. $10^9$	16 $10^5$

# A Fixed Target Experiment: A quarkonium observatory

- Interpolating the world data set:
- Rates expected at RHIC in 2011:  
 $J/\psi$ :  $10^6$  in  $pp$ ,  $\Upsilon$ :  $10^4$  in  $pp$
- 2-3 orders of magnitude higher here  
 (RHIC yields are much lower in dAu compared to pA here)
- Numbers are for only one unit of  $y$  about 0
- Unique access in the backward region
- Probe of the (very) large  $x$  in the target
- AIM/HOPE: Extract  $g(x, Q^2)$  with  $Q^2$  as low as  $10 \text{ GeV}^2$   
 from  $x = 10^{-3}$  up to  $\simeq \text{one}$

Target	$N_{J/\psi} (\gamma^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_{\Upsilon} (\gamma^{-1})$ <small><math>N_{\Upsilon} = A \mathcal{L} \sigma_{\Upsilon}</math></small>
(with branching and per unit of rapidity)		
Liq. H <sup>2</sup> (1m)	0.6 $10^9$	$10^6$
Liq. D <sup>2</sup>	1.5 $10^9$	23 $10^5$
Be	0.2 $10^9$	2.7 $10^5$
Cu	0.8 $10^9$	13 $10^5$
W	1.7 $10^9$	27 $10^5$
Pb	1. $10^9$	16 $10^5$



## A Fixed Target Experiment: A quarkonium observatory

- Interpolating the world data set:
- Rates expected at RHIC in 2011:  
 $J/\psi$ :  $10^6$  in  $pp$ ,  $\Upsilon$ :  $10^4$  in  $pp$
- 2-3 orders of magnitude higher here  
 (RHIC yields are much lower in dAu compared to pA here)
- Numbers are for only one unit of  $y$  about 0
- Unique access in the backward region
- Probe of the (very) large  $x$  in the target
- AIM/HOPE: Extract  $g(x, Q^2)$  with  $Q^2$  as low as  $10 \text{ GeV}^2$   
 from  $x = 10^{-3}$  up to  $\simeq \text{one}$
- Absolutely complementary to eRHIC (and LHeC)  
 whose focus would be low- $x$ .

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{J/\psi}</math></small>	$N_{\Upsilon} (y^{-1})$ <small><math>N_{\Upsilon} = A \mathcal{L} \sigma_{\Upsilon}</math></small>
(with branching and per unit of rapidity)		
Liq. H <sup>2</sup> (1m)	0.6 $10^9$	$10^6$
Liq. D <sup>2</sup>	1.5 $10^9$	23 $10^5$
Be	0.2 $10^9$	2.7 $10^5$
Cu	0.8 $10^9$	13 $10^5$
W	1.7 $10^9$	27 $10^5$
Pb	1. $10^9$	16 $10^5$



## A Fixed Target Experiment: A quarkonium observatory

- Interpolating the world data set:
- Rates expected at RHIC in 2011:  
 $J/\psi$ :  $10^6$  in  $pp$ ,  $\Upsilon$ :  $10^4$  in  $pp$
- 2-3 orders of magnitude higher here  
 (RHIC yields are much lower in dAu compared to pA here)
- Numbers are for only one unit of  $y$  about 0
- Unique access in the backward region
- Probe of the (very) large  $x$  in the target
- AIM/HOPE: Extract  $g(x, Q^2)$  with  $Q^2$  as low as  $10 \text{ GeV}^2$   
 from  $x = 10^{-3}$  up to  $\simeq \text{one}$
- Absolutely complementary to eRHIC (and LHeC)  
 whose focus would be low- $x$ .
- Use of  $pp$  vs  $pd$   $\rightarrow$  access to the  
 gluon content in the neutron in a wide  $x$  domain


Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{J/\psi}</math></small>	$N_{\Upsilon} (y^{-1})$ <small><math>N_{\Upsilon} = A \mathcal{L} \sigma_{\Upsilon}</math></small>
(with branching and per unit of rapidity)		
Liq. H <sup>2</sup> (1m)	0.6 $10^9$	$10^6$
Liq. D <sup>2</sup>	1.5 $10^9$	23 $10^5$
Be	0.2 $10^9$	2.7 $10^5$
Cu	0.8 $10^9$	13 $10^5$
W	1.7 $10^9$	27 $10^5$
Pb	1. $10^9$	16 $10^5$

cf. E866, Phys. Rev. Lett. 100 062301 (2008)

# A Fixed Target Experiment: A quarkonium observatory

Glue in the neutron and in the deuteron


E866 opened the way:  $g_n(x, Q^2 \simeq 100\text{GeV}^2) \simeq g_p(x, Q^2 \simeq 100\text{GeV}^2)$



# A Fixed Target Experiment: A quarkonium observatory

Glue in the neutron and in the deuteron

E866 opened the way:  $g_n(x, Q^2 \simeq 100\text{GeV}^2) \simeq g_p(x, Q^2 \simeq 100\text{GeV}^2)$




Such a measurement could be extended<sup>x2</sup>

# A Fixed Target Experiment: A quarkonium observatory

Glue in the neutron and in the deuteron

E866 opened the way:  $g_n(x, Q^2 \simeq 100\text{GeV}^2) \simeq g_p(x, Q^2 \simeq 100\text{GeV}^2)$




Such a measurement could be extended<sup>x2</sup>

- to lower  $x$ : down to  $\simeq 8 \times 10^{-3}$  with  $\Upsilon$

# A Fixed Target Experiment: A quarkonium observatory

Glue in the neutron and in the deuteron

E866 opened the way:  $g_n(x, Q^2 \simeq 100\text{GeV}^2) \simeq g_p(x, Q^2 \simeq 100\text{GeV}^2)$



Such a measurement could be extended<sup>x2</sup>


- to lower  $x$ : down to  $\simeq 8 \times 10^{-3}$  with  $\Upsilon$
- to lower  $Q^2$  with  $J/\psi$  (also to  $10\times$  lower  $x$ )

Beware of factorisation breaking effects for  $x_{projectile} \rightarrow 1$

# A Fixed Target Experiment: A quarkonium observatory

Glue in the neutron and in the deuteron

E866 opened the way:  $g_n(x, Q^2 \simeq 100\text{GeV}^2) \simeq g_p(x, Q^2 \simeq 100\text{GeV}^2)$



Such a measurement could be extended<sup>x2</sup>

- to lower  $x$ : down to  $\simeq 8 \times 10^{-3}$  with  $\Upsilon$
- to lower  $Q^2$  with  $J/\psi$  (also to  $10\times$  lower  $x$ )


Beware of factorisation breaking effects for  $x_{projectile} \rightarrow 1$

Quarkonium production on deuterium target could be analysed for  $x_{target} > 1$

# A Fixed Target Experiment: A quarkonium observatory

Glue in the neutron and in the deuteron

E866 opened the way:  $g_n(x, Q^2 \simeq 100\text{GeV}^2) \simeq g_p(x, Q^2 \simeq 100\text{GeV}^2)$



Such a measurement could be extended<sup>x2</sup>

- to lower  $x$ : down to  $\simeq 8 \times 10^{-3}$  with  $\Upsilon$
- to lower  $Q^2$  with  $J/\psi$  (also to  $10\times$  lower  $x$ )

Beware of factorisation breaking effects for  $x_{projectile} \rightarrow 1$


Quarkonium production on deuterium target could be analysed for  $x_{target} > 1$

- Need for high lumi: long deuterium target ?

# A Fixed Target Experiment: A quarkonium observatory

Glue in the neutron and in the deuteron

E866 opened the way:  $g_n(x, Q^2 \simeq 100\text{GeV}^2) \simeq g_p(x, Q^2 \simeq 100\text{GeV}^2)$



Such a measurement could be extended<sup>x2</sup>

- to lower  $x$ : down to  $\simeq 8 \times 10^{-3}$  with  $\Upsilon$
- to lower  $Q^2$  with  $J/\psi$  (also to  $10\times$  lower  $x$ )

Beware of factorisation breaking effects for  $x_{\text{projectile}} \rightarrow 1$

Quarkonium production on deuterium target could be analysed for  $x_{\text{target}} > 1$


- Need for high lumi: long deuterium target ?
- Unique probe of the deuteron internal dynamics (even for  $x_{\text{target}} < 1$ )



# A Fixed Target Experiment: A quarkonium observatory

Glue in the neutron and in the deuteron

E866 opened the way:  $g_n(x, Q^2 \simeq 100\text{GeV}^2) \simeq g_p(x, Q^2 \simeq 100\text{GeV}^2)$



Such a measurement could be extended<sup>x2</sup>

- to lower  $x$ : down to  $\simeq 8 \times 10^{-3}$  with  $\Upsilon$
- to lower  $Q^2$  with  $J/\psi$  (also to  $10\times$  lower  $x$ )

Beware of factorisation breaking effects for  $x_{\text{projectile}} \rightarrow 1$

Quarkonium production on deuterium target could be analysed for  $x_{\text{target}} > 1$

- Need for high lumi: long deuterium target ?
- Unique probe of the deuteron internal dynamics (even for  $x_{\text{target}} < 1$ )
- Momentum distribution of these gluons “shared” between  $n$  and  $p$  ?

# More on quarkonium as a probe...

- **New observables** involved quarkonium are needed to pin down the production mechanism see e.g. JPL, talk at Quarkonium Production, Vienna, 18-21 April 2001

# More on quarkonium as a probe...

- **New observables** involved quarkonium are needed to pin down the production mechanism      see e.g. JPL, talk at Quarkonium Production, Vienna, 18-21 April 2001
- They can also be promoted to **new probes**:

# More on quarkonium as a probe...

- **New observables** involved quarkonium are needed to pin down the production mechanism      see e.g. JPL, talk at Quarkonium Production, Vienna, 18-21 April 2001
- They can also be promoted to **new probes**:

## **Double $J/\psi$ production: a probe of gluon polarization?**

S.P. Baranov<sup>1</sup>, H. Jung<sup>2</sup>

<sup>1</sup>P.N.Lebedev Physical Institute, Moscow 117924, Russia

<sup>2</sup>III. Physikalisches Institut, Lehrstuhl B, RWTH Aachen, Germany

Received: 5 July 1994/Revised version: 5 October 1994      Z. Phys. C 66, 647–651 (1995)

**Abstract.** We consider the process of direct simultaneous production of two  $J/\psi$  particles and discuss the possibility that it can be used as a tool to measure the gluon polarization in the colliding particles.

# More on quarkonium as a probe...

- **New observables** involved quarkonium are needed to pin down the production mechanism      see e.g. JPL, talk at Quarkonium Production, Vienna, 18-21 April 2001
- They can also be promoted to **new probes**:

PHYSICAL REVIEW D

VOLUME 49, NUMBER 9

1 MAY 1994

## Associated $J/\psi + \gamma$ production as a probe of the polarized gluon distribution

M. A. Doncheski\*

*Department of Physics, University of Wisconsin, Madison, Wisconsin 53706*

C. S. Kim

*Department of Physics, Yonsei University, Seoul 120, Korea*

(Received 15 March 1993)

Associated production of  $J/\psi$  and a  $\gamma$  has recently been proposed as a clean probe of the gluon distribution. The same mechanism can be used to probe the polarized gluon content of the proton in polarized proton-proton collisions. We study  $J/\psi + \gamma$  production at both polarized fixed target and polarized collider energies.

# More on quarkonium as a probe...

- **New observables** involved quarkonium are needed to pin down the production mechanism see e.g. JPL, talk at Quarkonium Production, Vienna, 18-21 April 2001

- They can also be promoted to **new probes**:

Pair production of  $J/\psi$  as a probe of double parton scattering at LHCb

C. H. Kom\* and W. J. Stirling<sup>†</sup>

*Cavendish Laboratory, J.J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom*

A. Kulesza<sup>‡</sup>

*Institute for Theoretical Particle Physics and Cosmology,  
RWTH Aachen University D-52056 Aachen, Germany*

(Dated: May 24, 2011)

We argue that the recent LHCb observation of  $J/\psi$ -pair production indicates a significant contribution from double parton scattering, in addition to the standard single parton scattering component. We propose a method to measure the double parton scattering at LHCb using leptonic final states from the decay of two prompt  $J/\psi$  mesons.

# More on quarkonium as a probe...

- **New observables** involved quarkonium are needed to pin down the production mechanism see e.g. JPL, talk at Quarkonium Production, Vienna, 18-21 April 2001

- They can also be promoted to **new probes**:

Pair production of  $J/\psi$  as a probe of double parton scattering at LHCb

C. H. Kom\* and W. J. Stirling<sup>†</sup>

*Cavendish Laboratory, J.J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom*

A. Kulesza<sup>‡</sup>

*Institute for Theoretical Particle Physics and Cosmology,  
RWTH Aachen University D-52056 Aachen, Germany*

(Dated: May 24, 2011)

We argue that the recent LHCb observation of  $J/\psi$ -pair production indicates a significant contribution from double parton scattering, in addition to the standard single parton scattering component. We propose a method to measure the double parton scattering at LHCb using leptonic final states from the decay of two prompt  $J/\psi$  mesons.

- Double  $J/\psi$ ,  $J/\psi + \gamma$ ,  $J/\psi + D$ , ... can of course be studied with AFTER

## Part III

### AFTER as a quarkonium observatory in $pA$

(Precision analysis of Cold Nuclear Matter Effects)



# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable

# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**

# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**
- However, strong need for **cross checks** from various measurements

# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**
- However, strong need for **cross checks** from various measurements
  - **shadowing: open charm/beauty**

need for a vertex detector

# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**
- However, strong need for **cross checks** from various measurements
  - **shadowing: open charm/beauty**
  - **formation time:  $\chi_c, \eta_c, \psi'$**

need for a vertex detector

need for excellent photon calorimetry

# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**
- However, strong need for **cross checks** from various measurements
  - **shadowing: open charm/beauty**  
need for a vertex detector
  - **formation time:  $\chi_c, \eta_c, \psi'$**   
need for excellent photon calorimetry
  - **absorption: charmonium vs. bottomonium**  
need for luminosity

# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**
- However, strong need for **cross checks** from various measurements
  - **shadowing: open charm/beauty**  
need for a vertex detector
  - **formation time:  $\chi_c, \eta_c, \psi'$**   
need for excellent photon calorimetry
  - **absorption: charmonium vs. bottomonium**  
need for luminosity
- The **backward kinematics** is very useful for large- $x_{target}$  studies

# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**
- However, strong need for **cross checks** from various measurements
  - **shadowing: open charm/beauty**  
need for a vertex detector
  - **formation time:  $\chi_c, \eta_c, \psi'$**   
need for excellent photon calorimetry
  - **absorption: charmonium vs. bottomonium**  
need for luminosity
- The **backward kinematics** is very useful for large- $x_{target}$  studies
  - What is the amount of Intrinsic charm ? Is it color filtered ?  
relevant to understand E866 data



# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**
- However, strong need for **cross checks** from various measurements
  - **shadowing: open charm/beauty**  
need for a vertex detector
  - **formation time:  $\chi_c, \eta_c, \psi'$**   
need for excellent photon calorimetry
  - **absorption: charmonium vs. bottomonium**  
need for luminosity
- The **backward kinematics** is very useful for large- $x_{target}$  studies
  - What is the amount of Intrinsic charm ? Is it color filtered ?  
relevant to understand E866 data
  - **Is there an EMC effect for gluon ?** (reminder: EMC region  $0.3 < x < 0.7$ )

# A quarkonium observatory in $pA$ collisions

- The **target versatility** of a fixed-target experiment is undisputable
- A **wide rapidity coverage** is needed for:
  - a precise analysis of gluon nuclear PDF:  $y, p_T \leftrightarrow x_2$
  - a handle on **formation time effects** / the **energy dependence of  $\sigma^{abs}$**
- However, strong need for **cross checks** from various measurements
  - **shadowing: open charm/beauty**  
need for a vertex detector
  - **formation time:  $\chi_c, \eta_c, \psi'$**   
need for excellent photon calorimetry
  - **absorption: charmonium vs. bottomonium**  
need for luminosity
- The **backward kinematics** is very useful for large- $x_{target}$  studies
  - What is the amount of Intrinsic charm ? Is it color filtered ?  
relevant to understand E866 data
  - **Is there an EMC effect for gluon ?** (reminder: EMC region  $0.3 < x < 0.7$ )
  - In general one should be careful with factorization breaking effects:  
**This calls for different measurements to (in)validate factorization**

# A quarkonium observatory in $pA$ collisions

- Reminder:

Target	$N_{J/\Psi} (\gamma^{-1})$ <small><math>N_{J/\Psi} = A \mathcal{L} \sigma_{\Psi}</math></small>	$N_Y (\gamma^{-1})$ <small><math>N_Y = A \mathcal{L} \sigma_Y</math></small>
	<b>(with branching and per unit of rapidity)</b>	
Liq. H <sup>2</sup> (1m)	0.6 10 <sup>9</sup>	10 <sup>6</sup>
Liq. D <sup>2</sup>	1.5 10 <sup>9</sup>	23 10 <sup>5</sup>
Be	0.2 10 <sup>9</sup>	2.7 10 <sup>5</sup>
Cu	0.8 10 <sup>9</sup>	13 10 <sup>5</sup>
W	1.7 10 <sup>9</sup>	27 10 <sup>5</sup>
Pb	1. 10 <sup>9</sup>	16 10 <sup>5</sup>

# A quarkonium observatory in $pA$ collisions

- Reminder:
- Total yield measured by PHENIX during  $dAu$  Run08:  $9 \times 10^5 J/\psi$  (inclusive yield in nearly 3 units of  $y$ !)

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_Y (y^{-1})$ <small><math>N_Y = A \mathcal{L} \sigma_Y</math></small>
	(with branching and per unit of rapidity)	
Liq. $H^2$ (1m)	$0.6 \cdot 10^9$	$10^6$
Liq. $D^2$	$1.5 \cdot 10^9$	$23 \cdot 10^5$
Be	$0.2 \cdot 10^9$	$2.7 \cdot 10^5$
Cu	$0.8 \cdot 10^9$	$13 \cdot 10^5$
W	$1.7 \cdot 10^9$	$27 \cdot 10^5$
Pb	$1 \cdot 10^9$	$16 \cdot 10^5$

# A quarkonium observatory in $pA$ collisions

- Reminder:
- Total yield measured by PHENIX during  $dAu$  Run08:  $9 \times 10^5 J/\psi$  (inclusive yield in nearly 3 units of  $y$ !)
- Future plan for  $dAu$  runs at RHIC ?

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_Y (y^{-1})$ <small><math>N_Y = A \mathcal{L} \sigma_Y</math></small>
	(with branching and per unit of rapidity)	
Liq. $H^2$ (1m)	$0.6 \cdot 10^9$	$10^6$
Liq. $D^2$	$1.5 \cdot 10^9$	$23 \cdot 10^5$
Be	$0.2 \cdot 10^9$	$2.7 \cdot 10^5$
Cu	$0.8 \cdot 10^9$	$13 \cdot 10^5$
W	$1.7 \cdot 10^9$	$27 \cdot 10^5$
Pb	$1 \cdot 10^9$	$16 \cdot 10^5$

# A quarkonium observatory in $pA$ collisions

- Reminder:
- Total yield measured by PHENIX during  $dAu$  Run08:  $9 \times 10^5 J/\psi$  (inclusive yield in nearly 3 units of  $y$ !)
- Future plan for  $dAu$  runs at RHIC ?
- In principle, one can get **1000 times more  $J/\psi$**  (in 1 unit of  $y$ ), allowing for

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_Y (y^{-1})$ <small><math>N_Y = A \mathcal{L} \sigma_Y</math></small>
<b>(with branching and per unit of rapidity)</b>		
Liq. $H^2$ (1m)	<b>0.6 <math>10^9</math></b>	<b><math>10^6</math></b>
Liq. $D^2$	<b>1.5 <math>10^9</math></b>	<b>23 <math>10^5</math></b>
Be	<b>0.2 <math>10^9</math></b>	<b>2.7 <math>10^5</math></b>
Cu	<b>0.8 <math>10^9</math></b>	<b>13 <math>10^5</math></b>
W	<b>1.7 <math>10^9</math></b>	<b>27 <math>10^5</math></b>
Pb	<b>1. <math>10^9</math></b>	<b>16 <math>10^5</math></b>

# A quarkonium observatory in $pA$ collisions

- Reminder:
- Total yield measured by PHENIX during  $dAu$  Run08:  $9 \times 10^5 J/\psi$  (inclusive yield in nearly 3 units of  $y$ !)
- Future plan for  $dAu$  runs at RHIC ?
- In principle, one can get **1000 times more  $J/\psi$**  (in 1 unit of  $y$ ), allowing for
  - $\chi_c$  measurement in  $pA$  via  $J/\psi + \gamma$

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_{\gamma} (y^{-1})$ <small><math>N_{\gamma} = A \mathcal{L} \sigma_{\gamma}</math></small>
<b>(with branching and per unit of rapidity)</b>		
Liq. $H^2$ (1m)	<b>0.6 <math>10^9</math></b>	<b><math>10^6</math></b>
Liq. $D^2$	<b>1.5 <math>10^9</math></b>	<b>23 <math>10^5</math></b>
Be	<b>0.2 <math>10^9</math></b>	<b>2.7 <math>10^5</math></b>
Cu	<b>0.8 <math>10^9</math></b>	<b>13 <math>10^5</math></b>
W	<b>1.7 <math>10^9</math></b>	<b>27 <math>10^5</math></b>
Pb	<b>1. <math>10^9</math></b>	<b>16 <math>10^5</math></b>

# A quarkonium observatory in $pA$ collisions

- Reminder:
- Total yield measured by PHENIX during  $dAu$  Run08:  $9 \times 10^5 J/\psi$  (inclusive yield in nearly 3 units of  $y$ !)
- Future plan for  $dAu$  runs at RHIC ?
- In principle, one can get **1000 times more  $J/\psi$**  (in 1 unit of  $y$ ), allowing for
  - $\chi_c$  measurement in  $pA$  via  $J/\psi + \gamma$
  - **Polarisation** measurement as **function of  $A$ , the centrality,  $y$  and  $P_T$**  :  
For  $\alpha^{octet} \neq \alpha^{singlet}$ , probe of different absorption of octets & singlets ?

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_Y (y^{-1})$ <small><math>N_Y = A \mathcal{L} \sigma_Y</math></small>
<b>(with branching and per unit of rapidity)</b>		
Liq. $H^2$ (1m)	<b>0.6 <math>10^9</math></b>	<b><math>10^6</math></b>
Liq. $D^2$	<b>1.5 <math>10^9</math></b>	<b>23 <math>10^5</math></b>
Be	<b>0.2 <math>10^9</math></b>	<b>2.7 <math>10^5</math></b>
Cu	<b>0.8 <math>10^9</math></b>	<b>13 <math>10^5</math></b>
W	<b>1.7 <math>10^9</math></b>	<b>27 <math>10^5</math></b>
Pb	<b>1. <math>10^9</math></b>	<b>16 <math>10^5</math></b>



# A quarkonium observatory in $pA$ collisions

- Reminder:
- Total yield measured by PHENIX during  $dAu$  Run08:  $9 \times 10^5 J/\psi$  (inclusive yield in nearly 3 units of  $y$ !)
- Future plan for  $dAu$  runs at RHIC ?
- In principle, one can get **1000 times more  $J/\psi$**  (in 1 unit of  $y$ ), allowing for

- $\chi_c$  measurement in  $pA$  via  $J/\psi + \gamma$
- **Polarisation** measurement as **function of  $A$ , the centrality,  $y$  and  $P_T$**  :  
For  $\alpha^{octet} \neq \alpha^{singlet}$ , probe of different absorption of octets & singlets ?
- Ratio  $\psi'$  over **direct**  $J/\psi$  measurement in  $pA$

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_Y (y^{-1})$ <small><math>N_Y = A \mathcal{L} \sigma_Y</math></small>
<b>(with branching and per unit of rapidity)</b>		
Liq. $H^2$ (1m)	<b>0.6 <math>10^9</math></b>	<b><math>10^6</math></b>
Liq. $D^2$	<b>1.5 <math>10^9</math></b>	<b>23 <math>10^5</math></b>
Be	<b>0.2 <math>10^9</math></b>	<b>2.7 <math>10^5</math></b>
Cu	<b>0.8 <math>10^9</math></b>	<b>13 <math>10^5</math></b>
W	<b>1.7 <math>10^9</math></b>	<b>27 <math>10^5</math></b>
Pb	<b>1. <math>10^9</math></b>	<b>16 <math>10^5</math></b>

# A quarkonium observatory in $pA$ collisions

- Reminder:
- Total yield measured by PHENIX during  $dAu$  Run08:  $9 \times 10^5 J/\psi$  (inclusive yield in nearly 3 units of  $y$ !)
- Future plan for  $dAu$  runs at RHIC ?
- In principle, one can get **1000 times more  $J/\psi$**  (in 1 unit of  $y$ ), allowing for

Target	$N_{J/\psi} (y^{-1})$ <small><math>N_{J/\psi} = A \mathcal{L} \sigma_{\psi}</math></small>	$N_Y (y^{-1})$ <small><math>N_Y = A \mathcal{L} \sigma_Y</math></small>
<b>(with branching and per unit of rapidity)</b>		
Liq. $H^2$ (1m)	<b>0.6 <math>10^9</math></b>	<b><math>10^6</math></b>
Liq. $D^2$	<b>1.5 <math>10^9</math></b>	<b>23 <math>10^5</math></b>
Be	<b>0.2 <math>10^9</math></b>	<b>2.7 <math>10^5</math></b>
Cu	<b>0.8 <math>10^9</math></b>	<b>13 <math>10^5</math></b>
W	<b>1.7 <math>10^9</math></b>	<b>27 <math>10^5</math></b>
Pb	<b>1. <math>10^9</math></b>	<b>16 <math>10^5</math></b>

- $\chi_c$  measurement in  $pA$  via  $J/\psi + \gamma$
- **Polarisation** measurement as **function of  $A$ , the centrality,  $y$  and  $P_T$**  :  
For  $\alpha^{octet} \neq \alpha^{singlet}$ , probe of different absorption of octets & singlets ?
- Ratio  $\psi'$  over **direct**  $J/\psi$  measurement in  $pA$
- not to mention ratio with open charm, Drell-Yan, etc ...

## Part IV

# AFTER as a quarkonium observatory in $PbA$ (the quest for sequential suppression)

## A Fixed Target Experiment: a quarkonium observatory in $PbA$

Observation of  $J/\psi$  sequential suppression **seems to be hindered** by

- the **Cold Nuclear Matter effects**: non trivial and  
... not well-known, after all

## A Fixed Target Experiment: a quarkonium observatory in $PbA$

Observation of  $J/\psi$  sequential suppression **seems to be hindered** by

- the **Cold Nuclear Matter effects**: non trivial and  
... not well-known, after all
- the difficulty to observe directly the **excited states**  
which would melt before the ground states
  - $\chi_c$  **never studied in AA collisions**
  - $\psi(2S)$  **not yet studied in AA collisions at RHIC and the LHC**

## A Fixed Target Experiment: a quarkonium observatory in $PbA$

Observation of  $J/\psi$  sequential suppression **seems to be hindered** by

- the **Cold Nuclear Matter effects**: non trivial and  
... not well-known, after all
- the difficulty to observe directly the **excited states**  
which would melt before the ground states
  - $\chi_c$  **never studied in AA** collisions
  - $\psi(2S)$  **not yet** studied in AA collisions **at RHIC and the LHC**
- the possibilities for  **$c\bar{c}$  recombination**
  - **Open charm** studies are **difficult** where recombination matters most  
i.e. at **low  $P_T$**
  - Only indirect indications –from the  $y$  and  $P_T$  dependence of  $R_{AA}$ –  
that recombination may be at work
  - CNM effects may show a non-trivial  $y$  and  $P_T$  dependence too !
  - not clear what  $v_2$  tells us

## A Fixed Target Experiment: a quarkonium observatory in $PbA$

- The excellent capabilities in  $pA$  should help
  - to reduce the CNM uncertainties
  - to measure their dependence in  $y$  and  $P_T$

Rough estimation of the yield:  $2 \times 10^7 J/\psi$ ,  $10^4 Y$  per year ( $10^6$  sec)

## A Fixed Target Experiment: a quarkonium observatory in $PbA$

- The excellent capabilities in  $pA$  should help
  - to reduce the CNM uncertainties
  - to measure their dependence in  $y$  and  $P_T$
- Even though recombination may not be large at 72 GeV:
  - Open charm may be well measured, via displaced  $e/\mu$  or  $D \rightarrow K\pi$   
a priori even at low  $P_T$  thanks to the boost

Rough estimation of the yield:  $2 \times 10^7 J/\psi$ ,  $10^4 Y$  per year ( $10^6$  sec)



## A Fixed Target Experiment: a quarkonium observatory in $PbA$

- The excellent capabilities in  $pA$  should help
  - to reduce the CNM uncertainties
  - to measure their dependence in  $y$  and  $P_T$
- Even though recombination may not be large at 72 GeV:
  - Open charm may be well measured, via displaced  $e/\mu$  or  $D \rightarrow K\pi$   
a priori even at low  $P_T$  thanks to the boost
- last but not least, excited states would be studied
  - $\psi(2S)$  thanks to the statistics and the resolution
  - $\chi_c$  thanks the excellent calorimetry in high-multiplicity environment  
cf. the CALICE detector using particle flow techniques
  - and maybe ... for the very first time the  $\eta_c$

Rough estimation of the yield:  $2 \times 10^7 J/\psi$ ,  $10^4 Y$  per year ( $10^6$  sec)

## A Fixed Target Experiment: a quarkonium observatory in $PbA$

- The excellent capabilities in  $pA$  should help
  - to reduce the CNM uncertainties
  - to measure their dependence in  $y$  and  $P_T$
- Even though recombination may not be large at 72 GeV:
  - Open charm may be well measured, via displaced  $e/\mu$  or  $D \rightarrow K\pi$   
a priori even at low  $P_T$  thanks to the boost
- last but not least, excited states would be studied
  - $\psi(2S)$  thanks to the statistics and the resolution
  - $\chi_c$  thanks the excellent calorimetry in high-multiplicity environment  
cf. the CALICE detector using particle flow techniques
  - and maybe ... for the very first time the  $\eta_c$
- As STAR people suggested, why not to look for gluon quenching  
in  $J/\psi$ +hadron correlations vs. centrality  
(I suspect that we need a good  $pA$  baseline)

Rough estimation of the yield:  $2 \times 10^7 J/\psi$ ,  $10^4 Y$  per year ( $10^6$  sec)

## Part V

# AFTER as a quarkonium observatory in polarised collisions

(the quest for gluon spin contributions)

# Spin Physics with A Fixed Target Experiment at the LHC

- A further undisputable property of fixed-target experiments is  
the possibility of polarising the target

see COMPASS, HERMES, CLAS, ...

# Spin Physics with A Fixed Target Experiment at the LHC

- A further undisputable property of fixed-target experiments is  
the possibility of polarising the target

see COMPASS, HERMES, CLAS, ...

- The polarisation can be longitudinal and transverse

# Spin Physics with A Fixed Target Experiment at the LHC

- A further undisputable property of fixed-target experiments is  
the possibility of polarising the target  
see COMPASS, HERMES, CLAS, ...
- The polarisation can be longitudinal and transverse
- Single Transverse Spin Asymmetries unravel the correlations  
between the parton  $k_T$  and the proton spin  
→ information on orbital motion of partons in the proton !

# Spin Physics with A Fixed Target Experiment at the LHC

- A further undisputable property of fixed-target experiments is  
the possibility of polarising the target  
see COMPASS, HERMES, CLAS, ...
- The polarisation can be longitudinal and transverse
- Single Transverse Spin Asymmetries unravel the correlations  
between the parton  $k_T$  and the proton spin  
→ information on orbital motion of partons in the proton !
- Double Longitudinal Spin Asymmetries allow for the extraction of  
polarised PDFs

# Spin Physics with A Fixed Target Experiment at the LHC

- A further undisputable property of fixed-target experiments is  
the possibility of polarising the target  
see COMPASS, HERMES, CLAS, ...
- The polarisation can be longitudinal and transverse
- Single Transverse Spin Asymmetries unravel the correlations  
between the parton  $k_T$  and the proton spin  
→ information on orbital motion of partons in the proton !
- Double Longitudinal Spin Asymmetries allow for the extraction of  
polarised PDFs
- Double Transverse Spin Asymmetries probe transversity



# Spin Physics with A Fixed Target Experiment at the LHC

- A further undisputable property of fixed-target experiments is  
the possibility of polarising the target  
see COMPASS, HERMES, CLAS, ...
- The polarisation can be longitudinal and transverse
- Single Transverse Spin Asymmetries unravel the correlations  
between the parton  $k_T$  and the proton spin  
→ information on orbital motion of partons in the proton !
- Double Longitudinal Spin Asymmetries allow for the extraction of  
polarised PDFs
- Double Transverse Spin Asymmetries probe transversity
- The beam may become transversely polarised during the crystal  
extraction

M. Ukkhano, Nucl. Instrum. Meth. A 582 (2007) 378.

→ to be experimentally checked ...

# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:

# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:  
→ (single) **polarised**  $p^\uparrow p$  collisions, e.g. **Single Spin Asymmetry**

# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:

→ (single) **polarised**  $p^\uparrow p$  collisions, e.g. **Single Spin Asymmetry**

“a **nonzero** transverse **SSA** generated by a gluon Sivers TMD would be an **evidence against large contributions from CO transition**”.

F. Yuan, PRD 78, 014024 (2008).

# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:

→ (single) **polarised**  $p^\uparrow p$  collisions, e.g. **Single Spin Asymmetry**

“a **nonzero** transverse **SSA** generated by a gluon Sivers TMD would be an **evidence against large contributions from CO transition**”.

- $SSA \neq 0$  in  $pp$ : indication for CSM (as seen below) F. Yuan, PRD 78, 014024 (2008).

# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:

→ (single) **polarised**  $p^\uparrow p$  collisions, e.g. **Single Spin Asymmetry**

“a **nonzero** transverse **SSA** generated by a gluon Sivers TMD would be an **evidence against large contributions from CO transition**”.

- $SSA \neq 0$  in  $pp$ : indication for CSM (as seen below) F. Yuan, PRD 78, 014024 (2008).
- $SSA \neq 0$  in  $ep$ : indication for COM

# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:

→ (single) **polarised**  $p^\uparrow p$  collisions, e.g. **Single Spin Asymmetry**

“a **nonzero** transverse **SSA** generated by a gluon Sivers TMD would be an **evidence against large contributions from CO transition**”.

- $SSA \neq 0$  in  $pp$ : indication for CSM (as seen below) F. Yuan, PRD 78, 014024 (2008).
- $SSA \neq 0$  in  $ep$ : indication for COM
- it comes from the (im)possibility of final state interferences


# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:

→ (single) **polarised**  $p^\uparrow p$  collisions, e.g. **Single Spin Asymmetry**

“a **nonzero** transverse **SSA** generated by a gluon Sivers TMD would be an **evidence against large contributions from CO transition**”.

- SSA  $\neq 0$  in  $pp$ : indication for CSM (as seen below) F. Yuan, PRD 78, 014024 (2008).
- SSA  $\neq 0$  in  $ep$ : indication for COM
- it comes from the (im)possibility of final state interferences



PHENIX, PRD 82, 112008 (2010)




# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:

→ (single) **polarised**  $p^\uparrow p$  collisions, e.g. **Single Spin Asymmetry**

“a **nonzero** transverse **SSA** generated by a gluon Sivers TMD would be an **evidence against large contributions from CO transition**”.

- $SSA \neq 0$  in  $pp$ : indication for CSM (as seen below) F. Yuan, PRD 78, 014024 (2008).
- $SSA \neq 0$  in  $ep$ : indication for COM
- it comes from the (im)possibility of final state interferences



PHENIX, PRD 82, 112008 (2010)

- At  $x_F > 0$ , the gluon from the  $p^\uparrow$  has a larger  $x_B$


# Transverse Single Spin Asymmetry and quarkonia

Information on the  $Q$  production mechanisms can also be obtained in:

→ (single) **polarised**  $p^\uparrow p$  collisions, e.g. **Single Spin Asymmetry**

“a **nonzero** transverse **SSA** generated by a gluon Sivers TMD would be an **evidence against large contributions from CO transition**”.

- SSA  $\neq 0$  in  $pp$ : indication for CSM (as seen below) F. Yuan, PRD 78, 014024 (2008).
- SSA  $\neq 0$  in  $ep$ : indication for COM
- it comes from the (im)possibility of final state interferences



PHENIX, PRD 82, 112008 (2010)

- At  $x_F > 0$ , the gluon from the  $p^\uparrow$  has a larger  $x_B$
- It knows more about the proton spin than at low  $x_B \rightarrow$  SSA grows

# Spin Asymmetries and quarkonia

- For now, such Transverse SSA can be used to discriminate between production mechanisms

# Spin Asymmetries and quarkonia

- For now, such Transverse SSA can be used to discriminate between production mechanisms
- The situation is likely to change **in the future**, allowing us to **measure gluon Sivers function from quarkonia ( $J/\psi$ ,  $\chi_c$ ,  $\Upsilon$ )**

# Spin Asymmetries and quarkonia

- For now, such Transverse SSA can be used to discriminate between production mechanisms
- The situation is likely to change **in the future**, allowing us to **measure gluon Sivers function from quarkonia ( $J/\psi$ ,  $\chi_c$ ,  $\Upsilon$ )**
- It remains to be investigated **how quarkonium polarisation can be used** to form DSA

Attempt in: J. L. Cortes, B. Pire, Phys. Rev. **D38**, 3586 (1988).

# Spin Asymmetries and quarkonia

- For now, such Transverse SSA can be used to discriminate between production mechanisms
- The situation is likely to change **in the future**, allowing us to **measure gluon Sivers function from quarkonia ( $J/\psi$ ,  $\chi_c$ ,  $\Upsilon$ )**
- It remains to be investigated **how quarkonium polarisation can be used** to form DSA

Attempt in: J. L. Cortes, B. Pire, Phys. Rev. **D38**, 3586 (1988).

- If the beam gets transversely polarised, gluon transversity may be accessed (?)

# Spin Asymmetries and quarkonia

- For now, such Transverse SSA can be used to discriminate between production mechanisms
- The situation is likely to change **in the future**, allowing us to **measure gluon Sivers function from quarkonia** ( $J/\psi$ ,  $\chi_c$ ,  $\Upsilon$ )
- It remains to be investigated **how quarkonium polarisation can be used** to form DSA

Attempt in: J. L. Cortes, B. Pire, Phys. Rev. **D38**, 3586 (1988).

- If the beam gets transversely polarised, gluon transversity may be accessed (?)
- Of course, transverse SSA can be studied in parallel for other mesons ( $D$ ,  $B$ , ...)

# Spin Asymmetries and quarkonia

- For now, such Transverse SSA can be used to discriminate between production mechanisms
- The situation is likely to change in the future, allowing us to measure gluon Sivers function from quarkonia ( $J/\psi$ ,  $\chi_c$ ,  $\Upsilon$ )
- It remains to be investigated how quarkonium polarisation can be used to form DSA

Attempt in: J. L. Cortes, B. Pire, Phys. Rev. **D38**, 3586 (1988).

- If the beam gets transversely polarised, gluon transversity may be accessed (?)
- Of course, transverse SSA can be studied in parallel for other mesons ( $D$ ,  $B$ , ...)
- In general, the backward region is the most favourable allowing for measurements in the large  $x$  region of the polarised nucleon



# Part VI


## More with AFTER

(Drell-Yan, jet and W/Z)

# A Fixed Target Experiment

## A dilepton observatory

→ Region in  $x$  probed by dilepton production as function of  $M_{\ell\ell}$




# A Fixed Target Experiment

## A dilepton observatory

→ Region in  $x$  probed by dilepton production as function of  $M_{\ell\ell}$

→ Above  $c\bar{c}$ :  $x \in [10^{-3}, 1]$

→ Above  $b\bar{b}$ :  $x \in [9 \times 10^{-3}, 1]$



# A Fixed Target Experiment


## A dilepton observatory

→ Region in  $x$  probed by dilepton production as function of  $M_{\ell\ell}$

→ Above  $c\bar{c}$ :  $x \in [10^{-3}, 1]$

→ Above  $b\bar{b}$ :  $x \in [9 \times 10^{-3}, 1]$

**Note:**  $x_{\text{target}} (\equiv x_2) > x_{\text{projectile}} (\equiv x_1)$   
“backward” region



# A Fixed Target Experiment

## A dilepton observatory

→ Region in  $x$  probed by dilepton production as function of  $M_{\ell\ell}$


→ Above  $c\bar{c}$ :  $x \in [10^{-3}, 1]$

→ Above  $b\bar{b}$ :  $x \in [9 \times 10^{-3}, 1]$

**Note:**  $x_{\text{target}} (\equiv x_2) > x_{\text{projectile}} (\equiv x_1)$   
“backward” region

→ sea-quark asymmetries  
via  $p$  and  $d$  studies

- at large(est)  $x$ : backward (“easy”)
- at small(est)  $x$ : forward (need to stop the (extracted) beam)



# A Fixed Target Experiment

## A dilepton observatory

⇒ Region in  $x$  probed by dilepton production as function of  $M_{\ell\ell}$


→ Above  $c\bar{c}$ :  $x \in [10^{-3}, 1]$

→ Above  $b\bar{b}$ :  $x \in [9 \times 10^{-3}, 1]$

**Note:**  $x_{\text{target}} (\equiv x_2) > x_{\text{projectile}} (\equiv x_1)$   
“backward” region

→ sea-quark asymmetries  
via  $p$  and  $d$  studies


- at large(est)  $x$ : backward (“easy”)
- at small(est)  $x$ : forward (need to stop the (extracted) beam)



⇒ To do: to look at the rates to see how competitive this will be


# Overall

## Fixed Target @ LHC



# Overall


## Fixed Target @ LHC






# Overall

## Fixed Target @ LHC




# Overall

## Fixed Target @ LHC



# Overall

## Fixed Target @ LHC



## Part VII

### AFTER as photon-proton collider

# A Fixed Target Experiment

One exotic illustration of the potentialities:


Ultra-peripheral collisions

# A Fixed Target Experiment

One exotic illustration of the potentialities:

Ultra-peripheral collisions

Inelastic photoproduction of  $J/\psi$  via UPC\*




# A Fixed Target Experiment

One exotic illustration of the potentialities:

Ultra-peripheral collisions

Inelastic photoproduction of  $J/\psi$  via UPC\*




Thanks to the boost:  $W_{\gamma+p}^{\text{max}}$  for a coherent photon emission ( $Z^2$  fact.)  
can be as high as 25 GeV !

# A Fixed Target Experiment



One exotic illustration of the potentialities:

Ultra-peripheral collisions

Inelastic photoproduction of  $J/\psi$  via UPC\*



Thanks to the boost:  $W_{\gamma+p}^{max}$  for a coherent photon emission ( $Z^2$  fact.)  
can be as high as 25 GeV !



**Disclaimer:** these numbers suppose a dedicated trigger and are preliminary



# A photon-proton collider at the LHC ?

- Rates for **Inelastic  $J/\psi$  photoproduction** are  
large enough to be measured

# A photon-proton collider at the LHC ?

- Rates for **Inelastic  $J/\psi$  photoproduction** are  
large enough to be measured
- True also for **diffractive  $J/\psi$  photoproduction**

# A photon-proton collider at the LHC ?

- Rates for **Inelastic  $J/\psi$  photoproduction** are large enough to be measured
- True also for **diffractive  $J/\psi$  photoproduction**
- Handle on gluons (not sure though that one can compete in some way with EICs)

Z. Phys. C 76, 231–239 (1997)

ZEITSCHRIFT  
FÜR PHYSIK C  
© Springer-Verlag 1997

## Diffractive $J/\psi$ photoproduction as a probe of the gluon density

M.G. Ryskin<sup>1</sup>, R.G. Roberts<sup>2</sup>, A.D. Martin<sup>3</sup>, E.M. Levin<sup>1,4</sup>

<sup>1</sup> Petersburg Nuclear Physics Institute, 188350, Gatchina, St. Petersburg, Russia

<sup>2</sup> Rutherford Appleton Laboratory, Chilton, OX11 0QX, UK

<sup>3</sup> Department of Physics, University of Durham, Durham, DH1 3LE, UK

<sup>4</sup> School of Astronomy and Physics, Raymond and Beverly Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv, Israel

Received: 12 November 1996 / Revised version: 13 January 1997

**Abstract.** We use perturbative QCD, beyond the leading  $\ln Q^2$  approximation, to show how measurements of diffractive  $J/\psi$  production at HERA can provide a sensitive probe of the gluon density of the proton at small values of Bjorken  $x$ . We estimate both the effect of the relativistic motion of the  $c$  and  $\bar{c}$  within the  $J/\psi$  and of the rescattering of the  $c\bar{c}$  quark pair on the proton. We find that the available data for diffractive  $J/\psi$  photoproduction can discriminate between the gluon distributions of the most recent sets of partons.

# Part VIII

## Conclusion and outlooks

# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments

# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam  $\rightarrow 5 \times 10^8$  protons per sec

# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam  $\rightarrow 5 \times 10^8$  protons per sec
- This allows for high luminosity  $pp$ ,  $pA$  and  $PbA$  collisions at  $\sqrt{s} = 115$  GeV and  $\sqrt{s_{NN}} = 72$  GeV

# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam  $\rightarrow 5 \times 10^8$  protons per sec
- This allows for high luminosity  $pp$ ,  $pA$  and  $PbA$  collisions at  $\sqrt{s} = 115$  GeV and  $\sqrt{s_{NN}} = 72$  GeV
- **Example:** precision quarkonium studies taking advantage of



# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam  $\rightarrow 5 \times 10^8$  protons per sec
- This allows for high luminosity  $pp$ ,  $pA$  and  $PbA$  collisions at  $\sqrt{s} = 115$  GeV and  $\sqrt{s_{NN}} = 72$  GeV
- **Example:** precision quarkonium studies taking advantage of
  - high luminosity (reach in  $y$ ,  $P_T$ , small BR channels)
  - target versatility (CNM effects, strongly limited at colliders)
  - modern detection techniques (e.g.  $\gamma$  detection with high multiplicity)

# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments
- Extracting a few per cent of the beam  $\rightarrow 5 \times 10^8$  protons per sec
- This allows for high luminosity  $pp$ ,  $pA$  and  $PbA$  collisions at  $\sqrt{s} = 115$  GeV and  $\sqrt{s_{NN}} = 72$  GeV
- **Example:** precision quarkonium studies taking advantage of
  - high luminosity (reach in  $y$ ,  $P_T$ , small BR channels)
  - target versatility (CNM effects, strongly limited at colliders)
  - modern detection techniques (e.g.  $\gamma$  detection with high multiplicity)
- This would likely prepare the ground for  $g(x, Q^2)$  extraction

# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments
- Extracting **a few per cent of the beam**  $\rightarrow 5 \times 10^8$  **protons per sec**
- This allows for **high luminosity**  $pp$ ,  $pA$  and  $PbA$  collisions at  $\sqrt{s} = 115$  GeV and  $\sqrt{s_{NN}} = 72$  GeV
- **Example:** **precision quarkonium studies** taking advantage of
  - high luminosity (reach in  $y$ ,  $P_T$ , small BR channels)
  - target versatility (CNM effects, strongly limited at colliders)
  - modern detection techniques (e.g.  $\gamma$  detection with high multiplicity)
- This would likely prepare the ground for  **$g(x, Q^2)$  extraction**
- A wealth of possible measurements:  
DY, Open  $b/c$ , jet correlation, UPC... (not mentioning secondary beams)

# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments
- Extracting **a few per cent of the beam**  $\rightarrow 5 \times 10^8$  **protons per sec**
- This allows for **high luminosity**  $pp$ ,  $pA$  and  $PbA$  collisions at  $\sqrt{s} = 115$  GeV and  $\sqrt{s_{NN}} = 72$  GeV
- **Example:** **precision quarkonium studies** taking advantage of
  - high luminosity (reach in  $y$ ,  $P_T$ , small BR channels)
  - target versatility (CNM effects, strongly limited at colliders)
  - modern detection techniques (e.g.  $\gamma$  detection with high multiplicity)
- This would likely prepare the ground for  **$g(x, Q^2)$  extraction**
- A wealth of possible measurements:  
DY, Open  $b/c$ , jet correlation, UPC... (not mentioning secondary beams)
- Planned LHC long shutdown ( $< 2020$  ?) could be used to install the extraction system

# Conclusion and outlooks

- Both  $p$  and  $Pb$  LHC beams can be extracted without disturbing the other experiments
- Extracting **a few per cent of the beam**  $\rightarrow 5 \times 10^8$  **protons per sec**
- This allows for **high luminosity**  $pp$ ,  $pA$  and  $PbA$  collisions at  $\sqrt{s} = 115$  GeV and  $\sqrt{s_{NN}} = 72$  GeV
- **Example: precision quarkonium studies** taking advantage of
  - high luminosity (reach in  $y$ ,  $P_T$ , small BR channels)
  - target versatility (CNM effects, strongly limited at colliders)
  - modern detection techniques (e.g.  $\gamma$  detection with high multiplicity)
- This would likely prepare the ground for  **$g(x, Q^2)$  extraction**
- A wealth of possible measurements:  
DY, Open  $b/c$ , jet correlation, UPC... (not mentioning secondary beams)
- Planned LHC long shutdown ( $< 2020$  ?) could be used to install the extraction system
- Very good **complementarity** with electron-ion programs